

$\bar{B}^0 \rightarrow \pi^+ X$ in the Standard Model

X.-G. He¹, C.-P. Kao¹, J.P. Ma² and S. Pakvasa³

¹*Department of Physics, National Taiwan University, Taipei*

²*Institute of Theoretical Physics, Academia Sinica, Beijing*

and

³*Department of Physics and Astronomy, University of Hawaii, Honolulu, Hawaii, HI 96822*

(June, 2002)

Abstract

In this paper we investigate the possibility of studying $B \rightarrow \pi$ form factor using the semi-inclusive decays $\bar{B}^0 \rightarrow \pi^+ + X_q$. In general $B \rightarrow PX$ semi-inclusive decays involve several hadronic parameters. But for $\bar{B}^0 \rightarrow \pi^+ X_q$ decays we find that in the factorization approximation, the only unknown hadronic parameters are the form factors $F_{0,1}^{B \rightarrow \pi}$. Therefore these form factors can be studied in $\bar{B}^0 \rightarrow \pi^+ X_q$ decays. Using theoretical model calculations for the form factors the branching ratios for $\bar{B}^0 \rightarrow \pi^+ X_d (\Delta S = 0)$ and $\bar{B}^0 \rightarrow \pi^+ X_s (\Delta S = -1)$, with the cut $E_\pi > 2.1$ GeV, are estimated to be in the ranges of $(3.1 \sim 4.9) \times 10^{-5} (F_1^{B \rightarrow \pi}(0)/0.33)^2$ and $(2.5 \sim 4.2) \times 10^{-5} (F_1^{B \rightarrow \pi}(0)/0.33)^2$, respectively, depending on the value of γ . The combined branching ratio for $\bar{B}^0 \rightarrow \pi^+ (X_d + X_s)$ is about $7.4 \times 10^{-5} (F_1^{B \rightarrow \pi}(0)/0.33)^2$ and is insensitive to γ . We also discuss CP asymmetries in these decay modes.

Typeset using REVTeX

I. INTRODUCTION

There have been considerable efforts to understand the properties of B decays. Recently several rare charmless hadronic B decays have been measured. These decays being rare are sensitive to CP violating parameter γ in the Standard Model (SM) and also new physics [1–6]. While most of the studies have concentrated on the exclusive B decay modes, there are also some studies for semi-inclusive decays [3–5], for example $B \rightarrow \eta' X$ have been studied in details experimentally [7]. At present there are several other multi-body rare B decays have been measured, such as $B \rightarrow \pi\pi K, KKK$ [8]. It can be expected that more $B \rightarrow PX$ decay modes, with P being a light meson, will be experimentally studied. Theoretically at the quark level the effective Hamiltonian for B decays is well understood in the SM. The quark-hadron duality allows one to have a good understanding for inclusive hadronic decays. But for exclusive and semi-inclusive decays there are more uncertainties. The major uncertainties come from our poor understanding of the long distance strong interaction dynamics.

In rare charmless hadronic exclusive two body B decays of the type $B \rightarrow PP$, the operators which induce such decays in the SM, to the lowest order, are four quark operators O_i . In the factorization approximation, the four quark operators are factorized into bi-quark operators

$$\begin{aligned} \langle P_1 P_1 | O_i | B \rangle &= \langle P_1 | j_1 | 0 \rangle \langle P_2 | j_2 | B \rangle + \langle P_1 | j'_1 | 0 \rangle \langle P_2 | j'_2 | B \rangle + (1 \rightarrow 2, 2 \rightarrow 1) \\ &+ \langle P_1 P_2 | j_1 | 0 \rangle \langle 0 | j_2 | B \rangle + \langle P_1 P_2 | j'_1 | 0 \rangle \langle 0 | j'_2 | B \rangle, \end{aligned} \quad (1)$$

where $j'_1 \times j'_2$ is the Fierz transformed form of $j_1 \times j_2$. The last two terms are referred as the annihilation contributions which are usually assumed to be small and are neglected. The bi-quark operators $\langle P_i | j_1^{(\prime)} | 0 \rangle$ and $\langle P_i | j_2^{(\prime)} | B \rangle$ can be related to P meson decay constants or $B \rightarrow P$ transition form factors. Several of the decay constants, such as f_π and f_K have been measured to good accuracy, but less is known about form factors. Present experimental measurements are consistent with model calculations. To have better determinations of CP violating parameter and possible new physics, more accurate determination of the form

factors is necessary.

Some theoretical studies for semi-inclusive charmless hadronic decay modes $B \rightarrow PX$ have been carried out before [3–5]. In the factorization approximation, the decay amplitude contains several terms,

$$A(B \rightarrow PX) = \langle X|j_1|0 \rangle \langle P|j_2|B \rangle + \langle P|j_1|0 \rangle \langle X|j_2|B \rangle \\ + \langle PX|j_1|0 \rangle \langle 0|j_2|B \rangle + (\text{Fierz transformed terms}). \quad (2)$$

Compared with exclusive decays, semi-inclusive decays involve fewer unknown hadronic parameters, therefore semi-inclusive decays may be better understood from theoretical point of view. Experimentally semi-inclusive decays may be more difficult to study because background. In order to make sure that the observed events are from rare charmless B decays, and other processes, such as $B \rightarrow D(D^*)X' \rightarrow PX''$, do not contaminate the direct rare charmless hadronic $B \rightarrow PX$ decays, one needs to make a cut on the P energy. It has been shown that with a cut of $E_P > 2.1$ GeV, most of the unwanted events can be eliminated [3]. The resulting events will have a small invariant mass M_X^2 . With the cut $E_P > 2.1$ GeV, $M_X^2 < 5.7$ GeV². Rare charmless hadronic B semi-inclusive decays can be studied experimentally and useful information can be obtained. In this paper we study a class of the semi-inclusive decay $\bar{B}^0 \rightarrow \pi^+ X_q$ with the emphasis on the possibility of using this decays to determine the form factor $F_{0,1}^{B \rightarrow \pi}$.

II. DECAY AMPLITUDES FOR $\bar{B}^0 \rightarrow \pi^+ X_Q$

From Eq. (2) we see that in general there are three types of terms in the factorization approximation for the semi-inclusive decays of the type $B \rightarrow PX$. Each of the terms involves different hadronic parameters, with the first, the second and the third terms being proportional to B to P transition form factor from $\langle P|j_2|B \rangle$, the P decay constant from $\langle P|j_1|0 \rangle$, and some other parameters from $\langle PX|j_1|0 \rangle$ and $\langle 0|j_2|B \rangle$, respectively. If all three terms in Eq. (2) contribute with the same order of magnitude to a process,

the accumulated uncertainties will be substantial due to uncertainties in all the hadronic parameters involved, especially in the form factors.

If one or two terms in Eq. (2) can be eliminated, one can have a better understanding of the processes involved. Indeed this can be achieved by an appropriate choice of the initial meson B and the final meson P . For example in $\bar{B}^0 \rightarrow (K^- X), (\pi^- X)$, the term proportional to the form factors does not appear [4], and allow a good theoretical prediction. We find that there is only one possible choice for P where the second term is eliminated. These are the decays $\bar{B}^0 \rightarrow \pi^+ X_q$. Here X_d and X_s indicate the states having $\Delta S = 0$ and -1 , respectively. These decays are directly related to the form factors $F_{0,1}^{B \rightarrow \pi}$. Therefore $\bar{B}^0 \rightarrow \pi^+ X_q$ can be used to study these form factors and to test model calculations. The form factors can also be studied in semi-leptonic $B \rightarrow l \bar{\nu}_l \pi$ decays, probably with a better accuracy compared with that determined from $\bar{B}^0 \rightarrow \pi^+ X_q$. However, the final states are different, one in the leptonic environment and the other in hadronic environment. They are complementary to each other.

In $\bar{B}^0 \rightarrow \pi^+ X_q$, the bi-quark operators can only be in the forms: $j_1 = \bar{q}\Gamma_1 u$ and $j_2 = \bar{u}\Gamma_2 b$ and therefore

$$A(\bar{B}^0 \rightarrow \pi^+ X_q) = \langle X_q | j_1 | 0 \rangle \langle \pi^+ | j_2 | \bar{B}^0 \rangle + \langle X_q \pi^+ | j'_1 | 0 \rangle \langle 0 | j'_2 | \bar{B}^0 \rangle. \quad (3)$$

The second term being annihilation type is sub-leading. We will neglect its contribution. Note that for $q = s$, the annihilation term is automatically zero.

We would like to point out that $\bar{B}^0 \rightarrow \pi^+ X_q$ is a many-body decay. It is different from two-body decays. There are, in fact, more ways of factorization for a many-body decay, such as $\langle X_1 | j_1 | 0 \rangle \langle X'_1 \pi^+ | j_2 | \bar{B}^0 \rangle$ and $\langle X_2 \pi^+ | j'_1 | 0 \rangle \langle X'_2 | j'_2 | \bar{B}^0 \rangle$ and , with $X_q = X_1 + X'_1 = X_2 + X'_2$. The two terms in Eq. (3) correspond to the cases: $\langle X'_1 | = \langle 0 |$, $\langle X'_2 | = \langle 0 |$, respectively. For $\bar{B}^0 \rightarrow \pi^+ X_q$ with a cut $E_\pi > 2.1$ GeV, the final state X_q has a small invariant mass. This is a quasi-two-body decay, with π^+ and X_q moving rapidly apart in opposite directions. The probability of forming the final state $\langle X'_1 \pi^+ |$ with $\langle X'_1 | \neq \langle 0 |$ is less than the probability of forming the simple final state $\langle \pi^+ |$. The contribution of the

configuration $\langle X_2\pi^+|j'_1|0 \rangle \langle X'_2|j'_2|B \rangle$ is dominated by $\langle X_q\pi^+|j'_1|0 \rangle \langle 0|j'_2|\bar{B}^0 \rangle$ if it is not zero. The cases with $|X'_1 \rangle$ and $|X'_2 \rangle$ not equal to $|0 \rangle$ are also higher order in α_s and therefore suppressed.

We are now ready to present the detailed calculations. The effective Hamiltonian for rare charmless hadronic B decays at the quark level is given by

$$H_{eff} = \frac{G_F}{\sqrt{2}} \left\{ V_{ub}V_{uq}^*(c_1O_1 + c_2O_2) - \sum_{i=u,c,t} \sum_{n=3}^{10} V_{ib}V_{iq}^*c_n^iO_n \right\}. \quad (4)$$

Here O_n are quark and gluon operators and are given by

$$\begin{aligned} O_1 &= (\bar{s}_i u_j)_{V-A} (\bar{u}_j b_i)_{V-A}, \quad O_2 = (\bar{s}_i u_i)_{V-A} (\bar{u}_j b_j)_{V-A}, \\ O_{3(5)} &= (\bar{s}_i b_i)_{V-A} \sum_{q'} (\bar{q}'_j q'_j)_{V-(+)A}, \quad O_{4(6)} = (\bar{s}_i b_j)_{V-A} \sum_{q'} (\bar{q}'_j q'_i)_{V-(+)A}, \\ O_{7(9)} &= \frac{3}{2} (\bar{s}_i b_i)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_j q'_j)_{V+(-)A}, \quad O_{8(10)} = \frac{3}{2} (\bar{s}_i b_j)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_j q'_i)_{V+(-)A}, \end{aligned} \quad (5)$$

where $(V \pm A)(V \pm A) = \gamma^\mu(1 \pm \gamma_5)\gamma_\mu(1 \pm \gamma_5)$, $q' = u, d, s, c, b$, $e_{q'}$ is the electric charge number of the q' quark, and i and j are color indices.

The Wilson coefficients c_n^i have been calculated in different schemes [9,10]. In this paper we will use consistently the scheme independent NDR results. The values of c_n at $\mu \approx m_b$ with the next-to-leading order (NLO) QCD corrections are given by [10]

$$\begin{aligned} c_1 &= -0.307, \quad c_2 = 1.147, \quad c_3^t = 0.017, \quad c_4^t = -0.037, \quad c_5^t = 0.010, \quad c_6^t = -0.045, \\ c_7^t &= -0.0017\alpha_{em}, \quad c_8^t = 0.052\alpha_{em}, \quad c_9^t = -1.37\alpha_{em}, \quad c_{10}^t = -0.282\alpha_{em}, \\ c_{3,5}^{u,c} &= -\frac{1}{N_c}c_{4,6}^{u,c} = \frac{1}{N_c}P_s^{u,c}, \quad c_{7,9}^{u,c} = P_e^{u,c}, \quad c_{8,10}^{u,c} = 0, \end{aligned} \quad (6)$$

where $N_c = 3$ is the number of colors and $\alpha_{em} = 1/137$ is the electromagnetic fine structure constant. The functions $P_{s,e}^i$ are given by

$$\begin{aligned} P_s^i &= \frac{\alpha_s}{8\pi}c_2\left(\frac{10}{8} + G(m_I, \mu, k^2)\right), \quad P_e^i = \frac{\alpha_{em}}{9\pi}(N_c c_1 + c_2)\left(\frac{10}{9} + G(m_i, \mu, k^2)\right), \\ G(m, \mu, k^2) &= 4 \int_0^1 x(1-x) \ln \frac{m^2 - x(1-x)k^2}{\mu^2} dx. \end{aligned} \quad (7)$$

We obtain the decay amplitudes in the factorization approximation as

$$A(\bar{B}^0 \rightarrow \pi^+ X_q) = [\alpha_q \bar{q} \gamma_\mu (1 - \gamma_5) u + \beta_q \bar{q} \gamma_\mu (1 + \gamma_5) u] [F_1(q^2)(p_B^\mu + p_\pi^\mu) + (F_0(q^2) - F_1(q^2)) \frac{m_B^2 - m_\pi^2}{q^2} q^\mu + \gamma_q \bar{q} (1 + \gamma_5) u \frac{m_B^2 - m_\pi^2}{m_b - m_u} F_0(q^2)], \quad (8)$$

where $q = p_B - p_\pi$, and

$$\begin{aligned} \alpha_q &= \frac{G_F}{\sqrt{2}} [V_{ub} V_{uq}^* (\frac{1}{N_c} c_1 + c_2 + \frac{1}{N_c} c_3^{tu} + c_4^{tu} + \frac{1}{N_c} c_9^{tu} + c_{10}^{tu}) \\ &\quad + V_{cb} V_{cq}^* (\frac{1}{N_c} c_3^{tc} + c_4^{tc} + \frac{1}{N_c} c_9^{tc} + c_{10}^{tc})], \\ \beta_q &= \frac{G_F}{\sqrt{2}} [V_{ub} V_{uq}^* (c_6^{tu} + c_8^{tu}) + V_{cb} V_{cq}^* (c_6^{tc} + c_8^{tc})], \\ \gamma_q &= \frac{G_F}{\sqrt{2}} [V_{ub} V_{uq}^* (c_5^{tu} + c_7^{tu}) + V_{cb} V_{cq}^* (c_5^{tc} + c_7^{tc})] (-\frac{2}{N_c}), \end{aligned} \quad (9)$$

where $c^{ij} = c^i - c^j$. The above coefficients depend on the momentum exchange k^2 . In the heavy b quark limit, $k^2 = m_B^2(1 - 2E_q/m_B)$.

III. BRANCHING RATIOS FOR $\bar{B}^0 \rightarrow \pi^+ X_Q$

From the decay amplitudes obtained in the previous section, we obtain the differential branching ratio,

$$\frac{d\Gamma}{dx dy} = \frac{m_B^5}{16\pi^3} [(|\alpha_q|^2 + |\beta_q|^2) F_1^2(1-x)(x+y-1) + \frac{1}{4} |\gamma_q|^2 F_0^2(1-x)], \quad (10)$$

where $y = 2E_q/m_B$ and $x = 2E_\pi/m_B$. The physical integration intervals are: $0 < x < 1$ and $1-x < y < 1$. The branching ratios with appropriate cut on the π^+ energy $E_\pi > E_{cut}$ are given by

$$\Gamma(E_\pi > E_{cut}) = \int_{2E_{cut}/m_B}^1 dx \int_{1-x}^1 dy \frac{d\Gamma}{dx dy}. \quad (11)$$

There are several calculations of the form factors with the value to be typically in the range of $0.3 \sim 0.4$ [11,12]. We will use $F_{0,1}^{B \rightarrow \pi}(0) = 0.33$ for illustration. For $E_\pi > 2.1$ GeV, the dependence on q^2 is small, and we will use a single pole form as an approximation. For the KM matrix elements we will use the following independent variables, $V_{us} = \lambda$, $V_{ub} = |V_{ub}| \exp(-i\gamma)$, $V_{cb} = A\lambda^2$, with $\lambda = 0.2196$, $A = 0.835$, $|V_{ub}| = 0.08|V_{cb}|$. The CP

violating phase γ it treated as a free parameter. The results for the branching ratios are shown in Figure 1. We see that the branching ratios for $\bar{B}^0 \rightarrow \pi^+ X_d$ and $\bar{B}^0 \rightarrow \pi^+ X_s$ in the ranges of $(3.1 \sim 4.9) \times 10^{-5}$ and $(2.5 \sim 4.2) \times 10^{-5}$ respectively. These can be reached at B factories.




FIG. 1. The branching ratios as a function of γ .

We now would like to point out an interesting prediction in the SM regarding rate differences. Due to the unitarity property of the KM matrix elements, $Im(V_{ub}V_{ud}^*V_{cd}V_{cb}^*) = -Im(V_{ub}V_{us}^*V_{cs}V_{cb}^*)$, the rate differences, $\Delta_d = \Gamma(\bar{B}^0 \rightarrow \pi^+ X_d) - \bar{\Gamma}(B^0 \rightarrow \pi^+ \bar{X}_d)$ and the corresponding rate difference Δ_s have the same magnitude but opposite sign. When the final states X_d and X_s are not distinguished, one would get zero for the asymmetry, A ,

$$A = \frac{\Gamma(\bar{B}^0 \rightarrow \pi^+(X_d + X_s)) - \bar{\Gamma}(B^0 \rightarrow \pi^-(\bar{X}_d + \bar{X}_s))}{\Gamma(\bar{B}^0 \rightarrow \pi^+(X_d + X_s)) + \bar{\Gamma}(B^0 \rightarrow \pi^-(\bar{X}_d + \bar{X}_s))}. \quad (12)$$

This can provide a test for the Standard Model.

We also studied CP asymmetries, with the energy cut $E_\pi > 2.1$ GeV, defined by

$$A_{CP} = \frac{\Gamma(\bar{B}^0 \rightarrow \pi^+ X_q) - \bar{\Gamma}(B^0 \rightarrow \pi^- \bar{X}_q)}{\Gamma(\bar{B}^0 \rightarrow \pi^+ X_q) + \bar{\Gamma}(B^0 \rightarrow \pi^- \bar{X}_q)}. \quad (13)$$

The results for A_{CP} are shown in Figure 2. Since $\Delta_d = -\Delta_s$, because $Br(\bar{B}^0 \rightarrow \pi^+ X_d)$ has a larger value than $Br(\bar{B}^0 \rightarrow \pi^+ X_s)$ it is easy to understand why the asymmetry A_{CP} for

$\bar{B}^0 \rightarrow \pi^+ X_s$ is larger than $\bar{B}^0 \rightarrow \pi^+ X_d$. The asymmetry in $\bar{B}^0 \rightarrow \pi^+ X_d(X_s)$ can be as large as 5% (6%).




FIG. 2. The CP asymmetries as a function of γ .

IV. DISCUSSIONS AND CONCLUSIONS

From discussions in the previous sections, it can be seen that the measurements of the branching ratios for $\bar{B}^0 \rightarrow \pi^+ X_q$ can obtain information about the form factors $F_{1,0}^{B \rightarrow \pi}$. If the form factor is known, the branching ratios can be predicted. The numerical values are obtained using $F_1(0) = F_0(0) = 0.33$. In general $F_1(q^2)$ and $F_0(q^2)$ have different dependence on q^2 , one would expect that several hadronic parameters are needed to specify the details. However since $F_1(0) = F_0(0)$ for small q^2 , such as the case with $E_\pi > 2.1$ GeV, $F_1^{B \rightarrow \pi}(q^2) \approx F_0^{B \rightarrow \pi}(q^2) \approx F_1^{B \rightarrow \pi}(0)$, the branching ratios to a good approximation are proportional to $F_1^2(0)$. The branching ratios obtained can be normalized as, $Br(F_1) = Br(F_1(0) = 0.33)(F_1(0)/0.33)^2$. We have checked numerically with another set of realistic form factors having different q^2 dependence in Ref. [12]. We indeed find that in the kinematic region we are interested, the results do not change very much.

We have argued previously that experimental measurements for $\bar{B}^0 \rightarrow \pi^+ X_d$, although difficult, can be carried out with an appropriate cut on the energy of the pion E_π .

Combining the measurements with $X_q = X_d$ and $X_q = X_s$ can further enhance the statistic significance. We find that the combined branching ratio for $\bar{B}^0 \rightarrow \pi^+(X_d + X_s)$ is $\sim 7.4 \times 10^{-5}(F_1^{B \rightarrow \pi}(0)/0.33)^2$ which is insensitive to the phase angle γ . This implies that even without a good determination of γ , one can have useful information about the form factors. The combined branching ratio also makes the task easier in that one does not need to distinguish whether the final states X_q contains different strange numbers -1 or 0 . To ensure that the π^+ is from $\bar{B}^0 \rightarrow \pi^+ X_d$, and not from $B^0 \rightarrow \pi^+ \bar{X}_u$, tagging on $\bar{B}^0$ or B^0 is necessary this can be carried out at B factories. Measurements of $\bar{B}^0 \rightarrow \pi^+ X_q$ are therefore possible.

There are several uncertainties involved. Some of the main uncertainties are from the KM matrix elements and the phase γ . At present the best fit value for γ is around 60° [2]. If one uses the modes $\bar{B}^0 \rightarrow \pi^+ X_d$ and $\bar{B}^0 \rightarrow \pi^+ X_s$ individually, one need a good knowledge of γ to obtain precise information about the form factors. However, since the combined branching ratio has a very weak dependence on γ , using the combined ratio to obtain information on the form factors does not require a precise determination of γ . We studied the sensitivity of the branching ratios on the magnitudes of the KM matrix elements. The largest one may come from the magnitude of V_{ub} . The branching ratio of $\bar{B}^0 \rightarrow \pi^+ X_d$ is almost proportional to $|V_{ub}|^2$ which can be easily rescaled. The branching ratio of $\bar{B}^0 \rightarrow \pi^+ X_s$ is less sensitive to $|V_{ub}|$. For an accurate determination of the form factors, a good knowledge of the magnitude of the KM matrix elements, especially V_{ub} is important.

In this paper we have studied a class of semi-inclusive charmless hadronic B decays $\bar{B}^0 \rightarrow \pi^+ + X_q$. We find that for these decays, in the factorization approximation, the only unknown hadronic parameters are the form factors $F_{0,1}^{B \rightarrow \pi}$. Accurate measurement of these decays can provide important information about form factors. Using theoretical model calculations for the form factors the branching ratios for $\bar{B}^0 \rightarrow \pi^+ X_d (\Delta S = 0)$ and $\bar{B}^0 \rightarrow \pi^+ X_s (\Delta S = -1)$, with the cut $E_\pi > 2.1$ GeV, are estimated to be in the ranges of $(3.1 \sim 4.9) \times 10^{-5}(F_1^{B \rightarrow \pi}(0)/0.33)^2$ and $(2.5 \sim 4.2) \times 10^{-5}(F_1^{B \rightarrow \pi}(0)/0.33)^2$ respectively, with the combined branching ratio to be $7.4 \times 10^{-5}(F_1^{B \rightarrow \pi}(0)/0.33)^2$ almost independent of

γ , which are within the reach of B factories. $\bar{B}^0 \rightarrow \pi^+ X_q$ can provide interesting information about the form factors. CP violating asymmetries in these decays can be studied; and with the current knowledge of the KM phase γ , we expect the asymmetries to be around 5%.

ACKNOWLEDGMENTS

This work was supported in part by the ROC NSC under grant number NSC 90-2112-M-002-014, by the ROC Ministry of Education Academic Excellence Project 89-N-FA01-1-4-3, and by the U.S. D.O.E. under grant DE-FG-03-94ER40833.

REFERENCES

- [1] H.-Y. Cheng and K.-C. Yang, Phys. Rev. **D62**, 054029(2000); G. Deshpande et al., Phys. Rev. Lett. **82**, 2240 (1999); X.-G. He, W.-S. Hou and K.-C. Yang, Phys. Rev. Lett. **81**, 5738 (1998). A. Ali, G. Kramer and C.-D. Lu, Phys. Rev. **D58**, 094009 (1998); G. Deshpande, X.-G. He and J. Traneptic, Phys. Lett. **B 345**, 547 (1995); Y.-Y. Keum, H.-n. Li and A.I. Sanda, Phys. Rev. **D63**, 054008(2001); Phys. Lett. **B504**, 6(2001); C.-D. Lu, K. Ukai and M.-Z. Yang, Phys. Rev. **D63**, 074009(2001); D.-S. Du, D.-S. Yang and G.-H. Zhu, Phys. Lett. **B 488**, 46 (2000); Phys. Lett. **B488**, 46(2000); T. Muta, A. Sugamoto, M.-Z. Yang and Y.-D. Yang, Phys. Rev. **D 62**, 094020 (2000); M.-Z. Yang and Y.-D. Yang, Phys. Rev. **D 62**, 114019 (2000); M. Beneke et al., Phys. Rev. Lett. **83**, 1914 (1999); Nucl. Phys. **B 591**, 313 (2000).
- [2] M. Neubert and J. Rosner, Phys. Lett. **B441**, 403(1998); Phys. Rev. Lett. **81**, 5074(1998); M. Gronau and J. Rosner, Phys. Rev. **D57**, 6843(1998); X.-G. He et al., Phys. Rev. **D64**, 034002(2001); X.-G. He, C.-L. Hsueh and J.-Q. Shi, Phys. Rev. Lett. **84**, 18(2000); A. Buras and R. Fleischer, Eur. Phys. J. **C16**, 97(2000).
- [3] A. Datta et al., Phys. Rev. **D 57**, 6829 (1998).
- [4] X.-G. He, C.-H. Jin and J.-P. Ma, Phys. Rev. **D64**, 014020(2001); C.S. Kim et al., eprint hep-ph/0203093.
- [5] N. G. Deshpande et al., Phys. Lett. **B 366**, 300 (1996); A. Kagan and A. Petrov, e-print hep-ph/9707354; D. Atwood and A. Soni, Phys. Lett. **B405**, 150(1997); W.-S. Hou and B. Tseng, Phys. Rev. Lett. **80**, 434(1998); D. Atwood and A. Soni, Phys. Rev. Lett. **81**, 3324 (1998); A. Datta, X.-G. He and S. Pakvasa, Phys. Lett. **B 419**, 369 (1998); X.-G. He, J.P. Ma and C.-Y. Wu, Phys. Rev. **D63**, 094004(2001);
- [6] D. Ghosh et al., e-print hep-ph/0111106; X.-G. He, J.-Y. Leou and J.-Q. Shi, Phys. Rev. **D64**, 094018(2001); C.-K. Chua and W.-S. Hou, Phys. Rev. Lett. **86**, 2728(2001); X.-G. He, W.-S. Hou and K.-C. Yang, Phys. Rev. Lett. **81**, 5738(1998); M. Ciuchini, E.

- Gabrielli and G. Giudice, Phys. Lett. **B388**, 353(1996); F. Gabbiani et al., Nucl. Phys. **B477**, 321(1996).
- [7] D. Asner et al., CLEO collaboration, CONF 97-13, EPS332; B. Aubert et al., BABAR collaboration, hep-ex/0109034;
- [8] K. Abe et al., e-print hep-ex/0201001.
- [9] G. Buchalla, A. Buras and M. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996); A. Buras, M. Jamin and M. Lautenbacher, Nucl. Phys. **B 400**, 75 (1993); M. Ciuchini et al., Nucl. Phys. **B 415**, 403 (1994).
- [10] N. Deshpande and X.-G. He, Phys. Lett. **B 336**, 471 (1994).
- [11] M. Wirbel, B. Stech and M. Bauer, Z. Phys. **C29**, 637(1985); M. Bauer, B. Stech and M. Wirbel, Z. Phys. **34**, 103(1987); M. Bauer and B. Stech, Phys. Lett. **B152**, 380(1985).
- [12] Patricia Ball, e-print hep-ph/9802394, JHEP 9809, 005(1998).